

## ASSOCIATION OF HYPERTENSION AND DIASTOLIC DYSFUNCTION WITH TYPE-2 DIABETES MELLITUS

S.M. Sohail Ashraf<sup>1</sup>, Fasia Basir<sup>2</sup>

### ABSTRACT

**Objective:** To evaluate the prevalence of hypertension and diastolic dysfunction (LVDD) with normal ejection fraction in type-2 diabetes mellitus.

**Method:** Two hundred twelve diabetic patients (98 females and 114 males) from January 2005 to June 2006 were enrolled in this study. Their ages ranged from 35 to 65 years with mean age of 55 years. Patients with Diabetes Mellitus Type-2, Hypertension: BP<sup>3</sup>140/90 (with or without medication) Body Mass Index<sup>3</sup>30, Waist-Hip ratio>0.90. Left Ventricular Diastolic Dysfunction (LVDD) and Serum Triglyceride and HDL Cholesterol were included.

**Result:** Out of 212 patients, 104(49%) were found to be hypertensive (both systolic and diastolic) and 12 patients (5.6%) were found to be suffering from isolated systolic hypertension. Hence total number of hypertensive patients was 116 (54.6%). Thirty two patients were found to be suffering from diastolic dysfunction.

**Conclusion:** Hypertension and Diabetes are interlinked and show how micro and macrovascular complications of diabetes are increased when two conditions occur together. The LVDD is much more prevalent than previously suggested in patients with type-2 diabetes mellitus. Hence LVDD is an early marker of diabetic cardiomyopathy.

**KEYWORDS:** Diabetes Mellitus Type-2, Hypertension, Diastolic dysfunction.

Pak J Med Sci May - June 2007 Vol. 23 No. 3 344-348

### INTRODUCTION

Diabetes Mellitus is a cardiovascular disease. Two out of three diabetic patients die of cardiovascular disease or its complications.<sup>1</sup> According to the international diabetic federation the number of diabetic patients worldwide was estimated as 150 million in 2000, a figure that is said to rise to 300 million in 2025.<sup>2</sup> The

implications are enormous, both in terms of personal suffering and cost to health-care systems, which are increasingly faced with treating the serious macro-vascular complications of diabetes mellitus.

Hypertension frequently co exists with diabetes, there is an increased prevalence of hypertension among diabetic patients<sup>3</sup> but there is also high propensity among hypertensive patients to develop type-2 diabetes.<sup>4,5</sup> When occurring together the two disease entities appear to aggravate one another worsening both the diabetes and cardiovascular end points.<sup>6</sup>

Data from UK Prospective Diabetes Study (UKPDS) revealed that every 10mm.Hg reduction in the level of systolic BP is associated with a nearly 12% lower incidence in myocardial infarction, down to a systolic BP level of <120mm.Hg.<sup>7</sup> Progressive lower diastolic BP also reduces CV risk progressively. The Hyper-

1. Dr. S.M. Sohail Ashraf DTCD,MCPS,MD  
Assistant professor of Medicine
  2. Dr. Fasia Basir FCPS  
Associate professor of Medicine
- 1-2: Ziauddin Medical University Hospital  
Kemari Campus  
Karachi - Pakistan.

#### Correspondence

Dr. S.M. Sohail Ashraf,  
D-87/1, Block # 7,  
Gulshan-e-Iqbal,  
Karachi - Pakistan.  
E-mail: dr.sohailashraf@gmail.com

\* Received for Publication: September 7, 2006

\* Accepted: December 27, 2006

tension Optimal Treatment (HOT) trial showed that number of major cardiovascular events dropped in line with increasing astringent target diastolic BP-target.<sup>8</sup> These findings are reflected in the BP-target for diabetic patients of < 130/80mm of Hg as now recommended by European Society of Hypertension,<sup>9</sup> JNC7 report<sup>10</sup> and American Diabetes Association.<sup>11</sup>

Hypertension is the most important preventable cause of premature death in various countries. The ASCOT study (AngloScandinavian Cardiac Outcomes Trial) demonstrated both control of blood pressure and reduction in risk of strokes, heart attacks and other related diseases such as diabetes.<sup>12</sup>

Regarding diastolic dysfunction the most common feature of diabetic heart is abnormal early left-ventricular diastolic filling, suggesting reduced compliance or prolonged relaxation<sup>13</sup> because diabetes affect diastolic functions before systolic function,<sup>14</sup> left ventricular diastolic dysfunction may be an early marker of diabetic cardiomyopathy. Hypertension is also associated with impaired diastolic filling.<sup>15</sup>

## PATIENTS AND METHODS

A total of 212 patients (98 females and 114 males) of Diabetes Mellitus, 35 to 65 years of age (mean age was 55 years), from Ziauddin Medical University Hospital and its consultation clinic (Kaemari Campus) were enrolled in this study during January 2005 to June 2006. Prevalence of Hypertension and left ventricular diastolic dysfunction was carried out by means of cross-sectional study and all the patients were selected on random sample basis. The inclusion criteria was: Diabetes Mellitus Type-2, Hypertension: BP  $\geq$ 140/90 mm Hg. (with or without medication) Body Mass Index<sup>30</sup> Waist-Hip ratio $>$ 0.90.

Diastolic Dysfunction (Normal Systolic function LV ejection fraction  $\geq$ 50% no segmental wall motion abnormalities and no evidence of significant coronary valvular, infiltrative pericardial or pulmonary disease) and serum Triglyceride and HDL Cholesterol.

Patient on insulin therapy, those with history of angina pectoris or myocardial

infarction, Diabetic complications like Nephropathy, Neuropathy, and Retinopathy as well as valvular and congenital heart disease were excluded.

BP was measured with a random zero mercury sphygmomanometer. The mean of 6 measurements (3 while lying and 3 while standing) of systolic and diastolic BP was used on different occasions. As regards the assessment for obesity, BMI was calculated as weight in Kgs. divided by the square of height in meters. Waist circumference was calculated as an average of 2 measurements taken after inspiration and expiration at the mid-point of the lowest rib and iliac crest.

Waist-Hip ratio was defined as waist-girth divided by the hip-circumference measured at greater trochanter. For assessment of blood sugar, HbA1c and dyslipidaemia, the patients were asked to fast 12 to 14 hours. Then the blood samples were analysed. The reading of fasting blood glucose level, serum triglyceride, HDL, LDL Cholesterol plasma glucose concentration were measured by a glucose oxidase method and HbA1C was measured by an affinity binding assay, serum cholesterol, triglyceride and HDL were analyzed enzymatically. The reading of fasting blood glucose level, serum triglyceride, HDL, LDL cholesterol and HbA1c were recorded. For the assessment of effect on the CV system, ECG, chest X-ray and echocardiography were carried out.

Echocardiography was recorded by means of a 2-dimensional, M-mode and Doppler method. Patients were examined on the left lateral decubitus position using standard parasternal, short axis and apical views. Left ventricular diastolic dysfunction (LVDD) was evaluated. All Doppler measurements were assessed at the end of expiration.

From the transmitted recording, the measurements were carried out. Peak E- velocity in cms/sec. (Peak early transmitted filling velocity during early diastole). Peak A velocity in cms/sec (peak transmitted Atrial filling Velocity during late diastole) and deceleration time in milliseconds. (Time elapsed between peak E-velocity and the point where the extrapolation of

the declaration slope of the E-velocity crosses the zero base line). No subject had echocardiographically detectable regional wall motion abnormalities and each subject had normal ejection fractions. All the cardiac valves were examined to rule out significant valvular disease.

## RESULT

A total of 212 patients were screened and diagnosed with DM Type-2, their mean fasting blood sugar level was 140mg/dl. On average basis HBA1C was proved to be >8. Their mean age was calculated as 55 years. Mean duration of DM was 10 years. Out of 212 patients, 104(49%) patients were found to be hypertensive (both systolic and diastolic) and 12(5.6%) patients were suffering from systolic hypertension. Thus there were 116 hypertensive patients (54.6%). Among them both systolic and diastolic hypertension, there were 58 males and 46 females. Among systolic hypertensive patients 7 were males and 5 were females.

Serum Cholesterol was increased by 220mg/dl (mean). Serum Triglyceride was found to be

Table-I: Characteristic of 212 diabetic patients

Age in years (mean)	55 SD±6
Male	114
Female	98
Hypertension	116(54.7%)
BMI (Kg/m <sup>2</sup> ) (mean)	30 SD±2.8
Fasting blood sugar (mean)	140m/dL SD±12.9
HbA1C% (mean)	8 SD±0.95
Blood cholesterol (mean)	220mg/dL SD±20.40
Blood triglycerides (mean)	210mg/dL SD±19.5
HDL (mean)	44mg/dL SD±7.8
Waist-hip ratio	0.92 SD±0.06

Note: SD (Standard deviation)

Table-II: Hypertension and diastolic dysfunction ratio in Diabetes Mellitus

Sex	Total No. of patients	Hypertension	Diastolic dysfunction
	n=212	n=116	n=32
Male	114	65 (57%)	20(17.5%)
Female	98	51 (52%)	12(12.2%)

P-Value = 0.468 =0.283

210mg/dl (mean). Overall Body Mass Index was found to be 30 and waist-hip ratio was 0.92.

Diastolic dysfunction was carried out by means of 2-D (2 Dimensional) colored Doppler, and M-Mode echocardiography was performed. Only 32(15%) patients had shown isolated diastolic dysfunction having ejection fraction  $\geq 50$ . There were no ECG changes in the heart. All chest X-rays of patients were normal, so patients had Diabetes Mellitus Type-2 hypertension and diastolic dysfunction.

Thirty two out of 212 patients showed diastolic dysfunction with normal ejection fraction; subjects with impaired relaxation showed lower E-wave velocity compared with subjects with normal diastolic function. A-wave velocity was higher in patients with impaired relaxation compared with patients with normal diastolic dysfunction. Hence E:A ratio showed an inverse proportion or less than one. Most cases of diastolic dysfunction with normal ejection fraction were found in elderly age groups. There was no correlation between the E:A ratio and lipid profile, E:A ratio and HbA1C, E:A ratio and fasting blood glucose, E:A ratio and fasting blood glucose, E:A ratio and left ventricular mass.

Table-III: Involvement of different age groups

Ages (yrs)	Diabetes Mellitus	Hypertension	Diastolic dysfunction
35-45	20	6	2
46-55	94	52	13
56-65	98	58	17
Total	212	116	32

## DISCUSSION

According to the WHO estimates, there were >5.2 million Diabetics in the year 2000 and it will be 13.9 million by the year 2030. Pakistan is 6<sup>th</sup> in the world ranking.<sup>16</sup> Hence according to the DCCT and UKPDS, poor HbA1C control will be progressively worsening complications. Every 1% decrease in mean HbA1C (%) yields significant reduction in complications.<sup>17</sup> Hypertension is a common finding in patients with type-2 diabetes mellitus, when present it is an ominous sign that double the already elevated risk of future cardiovascular events.

Lowering BP has repeatedly shown to benefit hypertensive diabetic patients in terms of both macro and micro vascular disease. The fact that anti-hypertensive treatment reduces diabetic and CV end points in diabetes underlines the necessity to integrate rigorous BP control with glycaemic control in the management of diabetes.

In this study 212 diabetes mellitus patients were selected, out of which 116 patients (54.6%) proved to be hypertensive. Among them 12 patients were those who had systolic hypertension. They belonged to  $\geq 60$  years of age. FRAMINGHAM study revealed that on long term follow-up stroke and CHD are respectively 4 and 5 times more common in patients with systolic hypertension.<sup>18</sup> Both JNC-VII and WHO: (International Society of Hypertension) sub-committee guidelines have strongly recommended treating systolic hypertension at 140mm Hg. or even lower in diabetics. This isolated systolic hypertension is due to arterial stiffness, which results in an increased PWV (pulse wave velocity) and therefore reflects high pulse pressure.

There is a doubling of cardiovascular events when hypertension and diabetes mellitus coexist.<sup>19</sup> With each disease patient has abnormalities in central and peripheral characteristics of cardiovascular structure and function that precedes the clinical manifestation of cardiovascular disease, including increased left ventricular mass and wall thickness, left ventricular diastolic filling abnormalities impaired

endothelial functions and increased arterial stiffness. The most common feature of the diabetic heart is abnormal early left ventricular diastolic filling, suggesting reduced compliance and prolonged relaxation.<sup>13</sup> Diabetes affects diastolic function before systolic function; left ventricular diastolic dysfunction (LVDD) may be an early marker of diabetic cardiomyopathy. Hypertension is also associated with impaired diastolic filling.<sup>20</sup>

In our study there were 32 cases of diastolic dysfunction with EF  $\geq 50\%$ . This study shows inverse EA ratio, prolonged relaxation of left ventricle during M-Mode, 2-D and Doppler echocardiography though the number is small but it signifies alarming situation of diabetic heart in adults without any myocardial ischemia, congestive cardiac failure and any valvular heart disease.

As the age increases, there is a decrease in EA ratio. This progression of LVDD with aging seems markedly accelerated by diabetes. Hence increasing numbers of cases are found in elderly people in this study. Therefore LVDD can occur in individuals with well-controlled diabetes and without vascular complication. This diabetic cardiomyopathy may happen due to arterial stiffness, inflammation, obesity and endothelial dysfunction.<sup>21</sup>

## CONCLUSION

The evidence described clearly shows how closely Diabetes and hypertension are interlinked and how micro and macro vascular complications of Diabetes increase when the two conditions co-exist. The LVDD is much more prevalent than previously thought in patients with type 2 Diabetes Mellitus who are free of clinically detectable heart disease. Hence this study unmasks the significant number of LVDD. For prevention one should adopt life style modifications to control blood sugar level and a Dietary Approach to Stop Hypertension (DASH) eating plans.<sup>22</sup> (diet rich in fruits, vegetables and low fat dietary products with a reduced content of saturated and total fat.)

## REFERENCES

1. American Heart Association Circulation 1999;100:1134-46.
2. International Diabetic Federation Prevalence Estimates. August 2003.
3. Sowers JR, Epstein M, Frohlich ED. Diabetes hypertension, and cardiovascular disease. An update. Hypertension. 2001;37:1053-9.
4. Gress TW, Nieto FJ, Shahar E. Hypertension and anti-hypertensive therapy as risk factors for type-2 diabetes mellitus. Atherosclerosis Risk in Communities Study. N Engl J Med 2000;342:905-12.
5. Stolk RP, Van Splunder IP, Schouten JS. High blood pressure and the incidence of non-insulin dependent diabetes mellitus. Findings in a 11.5 year follow up study in The Netherlands. Eur J Epidemiol 1993;9:134-9.
6. Stamler J, Vaccaro O, Neaton JD. Diabetes, other risk factors, and 12 years cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial. Diabetes Care 1993;16:434-44.
7. Alder AL, Stratton IM, Neil HA. Association of systolic Blood pressure with macro vascular complications of type-2 diabetes (UKPDS 36): prospective observational study. BMJ 2000;321:412-9.
8. Hansson L, Zanchetti A, Carruthers SG. Effects of intensive blood pressure lowering and low-dose aspirin in patients with hypertension. Principal results of the Hypertension Optimal Treatment (HOT) randomized trial. HOT study group. Lancet. 1998;1755-62.
9. Guidelines Committee. European Society of Hypertension. European Society of cardiology guidelines for the management of arterial hypertension. J Hypertension 2003;21:1011-53.
10. The Seventh Report on the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. The JNC7 Report. JAMA 2003;289:2560-72.
11. American Diabetes Association. Treatment of hypertension in adults with diabetes. Diabetes Care. 2002;25(Suppl 1):S71-S73.
12. Clinical Trial Update European Society of Cardiology in Stockholm: Angloscandinavian Cardiac outcome trial (ASCOT) 2005. The Heart Org. Ciber Essay.
13. Uusitupa MI, Mustonen JN, Airaksinen KE. Diabetes heart muscle disease. Ann Med. 1990;22:377-86.
14. Raev DC. An Echocardiographic study of young Type-I diabetic patient. Diabetes Care. 1994;17:633-9.
15. Missault LH, Duprez DA, Brandt AA, De Buyzere ML, Adang LT, Clement DL. Exercise performance and diastolic filling in essential hypertension. Blood press. 1993;2:284-88.
16. World Health Organization: WHO: Country and regional data: World Health Report 2004.
17. Stratton IM UKPDS (35) BMJ 2000;321:405-12.
18. Anonymous. The sixth report of the joint national committee on prevention, detection, evaluation and treatment of high blood pressure. JNC-VI Arch of Int Med 1997;157:2413-66.
19. Kelsey A, Kelley K.S. Tran 2V walking and visiting blood pressure in adult. Prev Med 2001;33:120-7.
20. Missault LH, Duprez DA, Breandt AA, Dc buyzere ML, Adang LT, Clement DL. Exercise performance and diastolic filling in essential hypertension. Blood Pressure 1993;2:284-8.
21. Kervy J, Steward, ED. Exercise, Training and Cardiovascular consequences of type II Diabetes Mellitus and Hypertension JAMA 2003;15(1):22-30.
22. Sack FM, SuetKet LP, Vollmer WM. Effect on blood pressure of reduced dietary Sodium and dietary approach to stop hypertension (DASH) diet. DASH Sodium Collaborative Research Group. N Eng J Med 2001;344:3-10.