

Metabolic Syndrome and its determinants in a sample of young Iranian children with obesity

Fatemeh Esfarjani¹, Marjan Khalafi², Fatemeh Mohammadi³,
Negar Zamani-Nour⁴, Roya Kelishadi⁵

ABSTRACT

Objective: Childhood obesity increases the risk of metabolic syndrome (MetS) both in childhood and adulthood. The present study aimed to determine the prevalence of MetS and its potential determinants in a sample of Iranian obese children.

Methodology: This cross-sectional study was conducted in 2011 among 150 obese children (body mass index \geq 95th percentile) with seven years of age. They were randomly selected from 9 health centers in three districts of the north Tehran. Trained nutritionists completed a socio-demographic questionnaire by interviewing parents, and conducted the physical examination. MetS was defined based on modified Adult treatment panel III criteria.

Results: The mean (SD) of weight, height, and BMI was 37.5 (6.3) kg, 127.2 (4.7) cm and 23.08 (2.9) kg/m², respectively. The prevalence of MetS was 13.4%, without significant difference in terms of gender. The most common component of MetS was abdominal obesity (79%). While 21.3% of children did not have any component of MetS, 42% of them had at least one component. Most children with MetS had a history of breastfeeding for less than six months. Waist circumference, systolic and diastolic blood pressure, fasting blood glucose, and triglyceride levels were higher in MetS compared to controls ($p < 0.05$). Logistic regression model revealed that children with birth weight of \leq 2500 gr. were at higher risk of MetS than children with a higher birth weight (OR=4.3; 95%CI: 1.1-9.7).

Conclusion: Primordial prevention of childhood obesity, and screening the components of MetS among obese children, should be considered as a health priority at individual and public levels. Prevention of low birth weight can have long-term impact on prevention of childhood obesity.

KEY WORDS: Metabolic Syndrome, Children, Obesity, Prevention.

doi: [http://dx.doi.org/10.12669/pjms.291\(Suppl\).3511](http://dx.doi.org/10.12669/pjms.291(Suppl).3511)

How to cite this:

Esfarjani F, Khalafi M, Mohammadi F, Zamani-Nour N, Kelishadi R. Metabolic Syndrome and its determinants in a sample of young Iranian children with obesity. *Pak J Med Sci* 2013;29(1)Suppl:253-257. doi: [http://dx.doi.org/10.12669/pjms.291\(Suppl\).3511](http://dx.doi.org/10.12669/pjms.291(Suppl).3511)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Fatemeh Esfarjani,
 2. Marjan Khalafi,
 3. Fatemeh Mohammadi,
 4. Negar Zamani-Nour,
 5. Roya Kelishadi,
Department of Pediatrics,
Child Growth and Development Research Center,
Isfahan University of Medical Sciences, Isfahan, Iran.
- 1-4: Department of Food and Nutrition Policy and Planning Research,
National Nutrition and Food Technology Research Institute,
Faculty of Nutrition Sciences and Food Technology,
Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Correspondence:

Roya Kelishadi,
Department of Pediatrics,
Child Growth and Development Research Center,
Isfahan University of Medical Sciences,
Isfahan, Iran.
E-mail: Kelishadi@med.mui.ac.ir

INTRODUCTION

Metabolic Syndrome (MetS) is a collection of co-existence of disorders as hyperglycemia, dyslipidemia and hypertension.^{1,2} This syndrome increases the risk of chronic non-communicable diseases (CNCDs),³ which are a global emerging public health issues.⁴ The main cause of MetS remains to be determined but it is believed that this syndrome can be caused by a complex interaction between genetic, metabolic and environmental factors⁵⁻⁷ Childhood obesity is one of the most important public health problem in developing and developed countries.⁸ Obesity in children will also increase the risk of metabolic

syndrome in adulthood.⁹ More than 50% of obese children become obese adults with MetS in the future.¹⁰

There are various definitions for MetS.^{7,11,12} MetS is a considerable health problem in pediatric age group, notably among those with excess weight, though its reported prevalence varies in several studies due to lack of uniform criteria.¹³ The highest prevalence of MetS among over weight and obese children and adolescents were reported in Turkey (41.8%)¹⁴ and the United State (39%).¹⁵ The first nationwide studies in Middle-eastern adults¹⁶ and children and adolescents revealed considerably high prevalence of MetS. Considering that "School-age" is the most appropriate time for acquiring knowledge and habits, and given the impact of preventable factors on obesity and MetS,¹⁷⁻¹⁸ The present study was conducted to evaluate metabolic syndrome and its determinants in a sample of young Iranian children with obesity.

METHODOLOGY

This cross sectional study was conducted in 2011 within the framework of the "Assessing the effect of lifestyle modification intervention in school-age obese Children"¹⁹ on one hundred fifty, children with body mass index (BMI) $\geq 95^{\text{th}}$ percentile²⁰ who were selected randomly from three districts of health center in Tehran, Iran. Parents completed a socio-demographic questionnaire including age, birth order, birth weight & height, history of breastfeeding and duration, mothers' age, marital status, parents' occupation and education level, number of household members, through interviewing by trained nutritionists.

The Ethics Committee of the National Nutrition and Food Technology Research Institute approved the study. At the initial screening visit, oral consent was obtained from children and written consent from parents. Those children with genetic syndromes, any chronic diseases or disability as well as those with history of chronic medication use or those on special diet were not included.

Anthropometric Measurements: A team of trained health professionals measured anthropometric indexes including weight, height, and waist circumference (WC). Weight was measured to the nearest 0.1 kg using a calibrated and certified portable digital scale (Beurer, Germany) with lightly dressed, barefoot, and empty pockets. Height was measured in a standing position, barefoot and using a portable height gauge (Seca,

Germany) with accuracy of 0.1 cm. Then BMI was calculated as weight (kg) divided by height squared (m^2). WC was measured with an accuracy of 0.1 cm at the smallest area between the edge of the lower chest and the iliac crest bone.²¹ Before the examination, the scale was calibrated and the mean of two measurements was recorded.

Blood Pressure²²: BP was measured using a manual standard mercury sphygmomanometer from the right arm, after 5-10 minutes rest in sitting position by a qualified nurse. After that, the average of two measurements was recorded at the observer's eye level.²³

Biochemical Analysis: For Blood sampling, children were asked to fast overnight for 12 hours on the day of analysis and were invited to attend in accompaniment of one of their parents to a health center near the fieldwork at National Nutrition and Food Technology Research Institute. For measuring serum glucose and lipid concentrations, 5 ml blood was taken from the cubital vein. The fasting blood glucose (FBG), triglycerides (TG) and high-density lipoprotein-cholesterol (HDL-C) concentration were analyzed on fresh blood, using a commercial kit based on the enzymatic methods (Pars Azmoon, Iran) with auto-analyzer (Selectra E, Vita lab, Netherlands). Validity and reliability of measurement instruments in each of the cases were evaluated. After collecting blood samples, the children were served healthy breakfast and a gift.

Definition of MetS: We defined MetS based on ATP III modified for the pediatric age group²⁴ i.e. having three or more of the followings: FBG ≥ 100 mg/dl,²⁵ Fasting TG ≥ 110 mg/dl; HDL-C < 40 mg/dl; WC $\geq 90^{\text{th}}$ percentile for age and sex, according to national references curve;²⁶ systolic BP (SBP) or diastolic BP (DBP) $\geq 90^{\text{th}}$ percentile for age and sex and height, from national reference cut-off points.²⁷




Fig.1: Frequency of obese children according to the number of metabolic syndrome components.

Statistical Analysis: After recheck and editing the data, all statistical analyses were performed using SPSS software package (version 16; SPSS Inc., Chicago, IL, USA). Chi-square and the Student t tests were used to compare the ratios and the means of the groups. Pearson correlation coefficients were used to assess relationship between independent variable. Multivariate logistic regression analysis was used to determine the potential determinants of the MetS. Statistical significance was established at p-value <0.05.

RESULTS

This study comprised of 150 obese children (52% girls) with a mean (SD) age of 6.65±0.83. The mean (SD) of weight, height and BMI were 37.5 (6.3) kg, 127/2 (4.7) cm, and 23.1 (2/9) kg/m², respectively. The prevalence of MetS was 13.4%, without significant difference between girls and boys (10.3% vs. 16.7%, respectively>0.05). As depicted in Fig.1, 21.3% of children did not have any MetS components, 42% had at least one component, and 13.4% fulfilled the criteria of MetS and there were no children who had all of components.

As presented in Table-I, the majority of children with MetS had a birth weight of lower than 2500(gr) and were breastfed for less than 6 months. Other characteristics such as parents occupation and education, as well as family history of obesity were not significantly different among these children (data not shown). When the variables were simultaneously assessed into the logistic regression models, the risk of MetS was higher in those children with lower birth weight (≤2500gr) than in those with higher birth weight (OR=4.3; 95%CI: 1.1-9.7). The corresponding figure was not significant for other variables.

Table-I: Characteristics of obese children with or without metabolic syndrome.

Index	Overall (n=150)	Non-MetS (n=130)	MetS (n=20)	P value
Sex				
Girl	78 (100)	70 (89.7)	8 (10.3)	NS
Boy	72 (100)	60 (83.3)	12 (16.7)	
Birth order				
1	85 (100)	73 (85.9)	12 (14.1)	NS
≥2	65 (100)	57 (87.7)	8 (12.3)	
Birth weight(g)				
≤ 2500	15 (100)	10 (66.7)	5 (33.3)	0.035*
> 4000	131 (100)	116 (88.5)	15 (11.5)	
History of breastfeeding				
Yes	138 (100)	120 (87)	18 (13)	NS
No	12 (100)	10 (83.3)	2 (16.7)	
Duration of breastfeeding (mo)				
>6	75 (100)	61 (81.3)	14 (18.7)	0.045*
≥ 6	75 (100)	69 (92)	6 (8)	
Number of household members				
>3	55 (100)	49 (89.1)	6 (10.9)	NS
≤4	95 (100)	81 (85.3)	14 (14.7)	
Mothers' age (year)				
≥32	41 (100)	34 (82.9)	7 (17.1)	NS
<32	108 (100)	95 (88)	13 (12)	
Gestational Diabetes				
Yes	8 (100)	7 (87.5)	1 (12.5)	NS
No	142 (100)	123 (86.6)	19 (13.4)	

Data are presented as number (percent) or mean (SD)
*p < 0.05 (significantly different between two groups);
NS: non-significant MetS: metabolic syndrome

The mean (SD) of metabolic risk factors in obese children with and without MetS is presented in Table-II. Waist circumference, systolic and diastolic blood pressure, fasting blood glucose, and triglyceride levels were higher in MetS compared

Table-II: Comparison of metabolic risk factors in obese children with and without metabolic syndrome.

Index	Overall (n=150)	Non-MetS (n=130)	MetS (n = 20)	P value
BMI (kg/m ²)	23.07 (2.90)	22.99 (2.99)	23.59 (2.24)	NS
WC (cm)	74.02 (6.55)	73.84 (6.68)	75.20 (5.66)	* 0.047
SBP (mmHg)	99.61 (11.16)	97.16 (8.93)	115.50 (11.39)	* 0.000
DBP (mmHg)	62.59 (8.54)	60.67 (6.58)	75.05 (9.42)	* 0.000
FBG (mg/ dl)	85.65 (8.31)	84.91 (7.72)	90.05 (10.61)	* 0.010
TG (mg/ dl)	103.53 (46.27)	99.58 (45.63)	129.15 (43.03)	* 0.007
HDL-C (mg/ dl)	51.07 (9.42)	51.33 (8.92)	49.30 (12.30)	NS

Data are presented as mean (±SD)

*p < 0.05 (significantly different between two groups); NS: non-significant

MetS: metabolic syndrome; BMI: body mass index; WC: waist circumferences; SBP: systolic blood pressure;

DBP: diastolic blood pressure; FBG: fasting blood glucose; TG: triglyceride;

HDL-C: high-density lipoprotein-cholesterol.

to controls ($p < 0.05$). Significant relationship was documented for WC and BMI ($r = 0.85$, $p < 0.001$), SBP ($r = 0.25$, $p < 0.01$), and DBP ($r = 0.22$, $p < 0.01$). The most common components of MetS were abdominal obesity (79%) followed by high serum TG (74%) and high DBP (74%). High FBG has the lowest percentage in those components (21%). The prevalence of MetS components was not significantly different in terms of gender.

DISCUSSION

The rapid rising prevalence of childhood obesity is related to increased risk of metabolic syndrome. In our study, the prevalence of metabolic syndrome in obese children according to ATP III criteria was similar to Brazil (13.4% vs. 13/2%)²⁸ and other reports²⁹⁻³⁵ from 10% to 20%. Taylor discussed in her review that the prevalence of MetS among children and adolescents around the world is on average about 10%, ranging from near 2% among normal weight children and adolescents to about 32% among the obese.³⁶ The range of discrepancies in the prevalence of MetS were considerable in various studies, and can be attributed to ununified definition of this syndrome, different cut-off values, population size, age and degree of obesity in children.

The role of breastfeeding in prevention of obesity and MetS was shown in previous studies.^{27,37-38} It has been reported that the prevalence of obesity were twice in children who were not breastfed as compared to breastfed children, with a concurrent increase in MetS prevalence.³⁹ There was no relationship between breastfeeding and metabolic syndrome in a study by Kelishadi et al,⁴⁰ although the duration of breastfeeding has protective effect against MetS. Birth weight reflects the pattern of intrauterine growth and might have long-term impacts on chronic disease in adulthood.⁴¹ Our results showed the significant correlation between MetS and birth weight and supported the reports in studies that demonstrate the effect of low birth weight on the development of MetS in adulthood.⁴²⁻⁴³ We also measured waist circumferences, as a good predictor of cardiovascular disease (CVD) risk factors and MetS in children and adolescents.⁴⁴⁻⁴⁵

Increased waist circumference indicates abdominal obesity that was considered an important factor for the other components of MetS.⁴⁶ The high prevalence of hypertriglyceridemia may

be explained by role of dietary habits, for instance diets high in Trans fatty acid. This was not evaluated in our survey. The children's high blood pressure in this study was in keeping with the existing knowledge that abdominal obesity is clearly linked to hypertension.^{47,48} In this study the prevalence of hyperglycemia was lowest among the components of metabolic syndrome, which is consistent with some previous studies.^{24,49} It is believed that hyperglycemia probably is increased later than the other components of metabolic syndrome and may be associated with the duration of obesity.

Considering the global problem of childhood obesity,⁵⁰ Detecting MetS at an early stage and improving it by life style changes should be considered as an important goal in childhood. Early primary prevention to reduce weight, increase physical activities and train the parents of obese children are necessary. It is recommended that lifestyle modifications and plans for weight loss should be at the core of treating and preventing MetS and its components.

CONCLUSION

Although the prevalence of MetS depends on the definition used and population studied, it has been clearly increasing globally in parallel with childhood obesity and it can be a warning for the policy priorities in health strategic plans. Need to be alerted regarding the existence of the metabolic syndrome in obese children can reduce the health care costs. The health politics, potential screening and mass intervention programs at early age groups are crucial. Establishment of a uniform and universally accepted set of criteria for defining overweight, obesity and the metabolic syndrome in children would be a foundation for addressing this emerging public health concern. Further national prospective studies with long-term follow up in children with different ages are needed.

ACKNOWLEDGEMENT

The authors would like to appreciate the Research Council of National Nutrition and Food Technology Research Institute, Faculty of Nutrition Science and Food Technology, Shahid Beheshti University of Medical Sciences for financial support. (Grant No. 2676) We would also like to thank all parents and children for their valuable help in conducting this study. In addition, we thank Dr. Arghavan Etebarian and Roshanak Roustae for their technical support.

REFERENCES

1. Cruz ML, Weigensberg MJ, Huang TTK, Ball G, Shaibi GQ, Goran MI. The metabolic syndrome in overweight Hispanic youth and the role of insulin sensitivity. *J Clin Endocrinol Metabol.* 2004;89(1):108-113.
2. Reaven GM. Banting lecture 1988. Role of insulin resistance in human disease. *Diabetes.* 1988;37(12):1595-1607.
3. Kelishadi R. Childhood overweight, Obesity, metabolic syndrome in developing Countries. *Epidemiol Reviews.* 2007;29(1):62-76.
4. Galal O. Nutrition-related health patterns in the Middle East. *Asia Pacific J Clin Nutr.* 2003;12(3):337.
5. Groop L. Genetics of the metabolic syndrome. *Br J Nutr.* 2000;83(1):39.
6. Kelishadi R, Poursafa P. Obesity & Air Pollution: Global Risk Factors for Pediatric Non-alcoholic Fatty Liver Disease. *Hepatitis Monthly.* 2011;11(10):794.
7. Lidfeldt J, Nyberg P, Nerbrand C, Ojehagen A, Samsioe G, Schersten B. Biological factors are more important than socio-demographic and psychosocial conditions in relation to hypertension in middle-aged women. The Women's Health in the Lund Area (WHILA) study. *Blood Pressure.* 2002;11(5):270-278.
8. Karnik S, Kanekar AS. Childhood Obesity: A Global Public Health Crisis. *Int J Prev Med.* 2011;3(1):1-7.
9. Park M, Falconer C, Viner R, Kinra S. The impact of childhood obesity on morbidity & mortality in adulthood: a systematic review. *Obes Rev.* 2012;13(11):985-1000.
10. Vanhala M, Vanhala P, Kumpusalo E, Halonen P, Takala J. Relation between obesity from childhood to adulthood & the metabolic syndrome: population based study. *BMJ.* 1998;317(7154):319-320.
11. Ford ES, Li C, Zhao G, Pearson WS, Mokdad AH. Prevalence of the metabolic syndrome among US adolescents using the definition from the International Diabetes Federation. *Diabetes Care.* 2008;31(3):587-589.
12. Morrison J, Ford E, Steinberger J. The pediatric metabolic syndrome. *Minerva Med.* 2008;99(3):269-287.
13. De Ferranti SD, Osganian SK. Epidemiology of paediatric metabolic syndrome & type 2 diabetes mellitus. *Diab Vasc Dis Res.* 2007;4(4):285-296.
14. Sen Y, Kandemir N, Alikasifoglu A, Gonc N, Ozon A. Prevalence & risk factors of metabolic syndrome in obese children & adolescents: the role of the severity of obesity. *Euro J Pediatr.* 2008;167(10):1183-1189.
15. Shaibi GQ, Goran MI. Examining metabolic syndrome definitions in overweight Hispanic youth: a focus on insulin resistance. *J Pediatr.* 2008;152(2):171-176.
16. Delavari A, Forouzanfar MH, Alikhani S, Sharifian A, Kelishadi R. First Nationwide Study of the Prevalence of the Metabolic Syndrome & Optimal Cutoff Points of Waist Circumference in the Middle East. *Diabetes Care.* 2009;32(6):1092-1097.
17. Esmaillzadeh A, Mirmiran P, Azadbakht L, Azizi F. Changes in overweight prevalence among Tehrani children and adolescents: comparison of three different definitions. *Razi J Med Sci.* 2006;13(52):19-30.
18. Kelishadi R, Ghatrehsamani S, Hosseini M, Mirmoghataee P, Mansouri S, Poursafa P. Barriers to Physical Activity in a Population-based Sample of Children and Adolescents in Isfahan, Iran. *Int J Prev Med.* 2010;1(2):131.
19. Esfarjani F. Assessing the effect of lifestyle modification intervention in school-age obese Children. Final Report: National Nutrition & Food Technology Research Institute. Tehran, Iran. 2012.
20. Kuczumarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R. CDC growth charts: United States. *Advance Data.* 20.1:(314).
21. Kelishadi R, Ardalan G, Gheiratmand R, Adeli K, Delavari A, Majdzadeh R. Paediatric metabolic syndrome and associated anthropometric indices: the CASPIAN Study. *Acta Paediatr.* 2006;95(12):1625-1634.
22. Gargari BP, Behzad MH, Ghassabpour S, Ayat A. Prevalence of overweight and obesity among high-school girls in Tabriz, Iran, in 2001. *Food & Nutr Bulletin.* 2004;25(3):288-291.
23. Group NHBPEPW. Update on the 1987 task force report on high blood pressure in children and adolescents: A working group report from the National High Blood Pressure Education Program. *Pediatr.* 1996;98(4):649-658.
24. Cook S, Weitzman M, Auinger P, Nguyen M, Dietz WH. Prevalence of a metabolic syndrome phenotype in adolescents: findings from the third National Health and Nutrition Examination Survey, 1988-1994. *Arch Pediatr Adolesc Med.* 2003;157(8):821-827.
25. American Diabetes Association: Diagnosis and classification Diabetes Mellitus (Position statement). *Diabetes Care* 2004;(Suppl 1):10-15.
26. Kelishadi R, Gouya MM, Ardalan G, Hosseini M, Motaghian M, Delavari A. First reference curves of waist and hip circumferences in an Asian population of youths: CASPIAN study. *J Tropical Pediatr.* 2007;53(3):158-164.
27. Kelishadi R, Ardalan G, Gheiratmand R, Majdzadeh R, Delavari A, Heshmat R. Blood pressure and its influencing factors in a national representative sample of Iranian children and adolescents: The CASPIAN Study. *Eur J Cardiovasc Prev Rehabil.* 2006;13(6):956-963.
28. Barzin M, Hosseinpanah F, Saber H, Sarbakhsh P, Nakhoda K, Azizi F. Gender Differences Time Trends for Metabolic Syndrome and Its Components among Teheranian Children and Adolescents. *Cholesterol.* 2012;2012.
29. Calcaterra V, Klersy C, Muratori T, Telli S, Caramagna C, Scaglia F. Prevalence of metabolic syndrome (MS) in children and adolescents with varying degrees of obesity. *Clin Endocrinol.* 2008;68(6):868-872.
30. Ferreira AP, Ferreira CB, Brito CJ, Pitanga FJG, Moraes CF, Naves LA. Prediction of metabolic syndrome in children through anthropometric indicators. *Arquivos Brasileiros de Cardiologia.* 2011;96(2):121-125.
31. Fu J, Liang L, Zou C, Hong F, Wang C, Wang X. Prevalence of the metabolic syndrome in Zhejiang Chinese obese children and adolescents and the effect of metformin combined with lifestyle intervention. *Int J Obes.* 2006;31(1):15-22.
32. Mimoun E, Aggoun Y, Pousset M, Dubern B, Bougle D, Girardet JP. Association of arterial stiffness and endothelial dysfunction with metabolic syndrome in obese children. *J Pediatr.* 2008;153(1):65-70.
33. Tapia Ceballos L, Lpez Siguero JP, Jurado Ortiz A. [Prevalence of metabolic syndrome and its components in obese children and adolescents]. *An Pediatr (Barc).* 2007 Oct;67(4):352-61. Spanish
34. Ebrahimpour P, Pourebrahim R, Hamidi A, Fakhrazadeh D, Heshmat R, Nouri M. Metabolic Syndrome and its relationship with insulin levels in obese children: A study on primary school children of the 6th districts of Tehran. *J Iran Uni Med Sci.* 2006;13(51):7-15.
35. Ghaemi N, Aghaei Monavar A. Prevalence of Metabolic Syndrome in children with overweight. *Med J Mashhad Uni Med Sci.* 2010.
36. Tailor AM, Peeters PHM, Norat T, Vineis P, Romaguera D. An update on the prevalence of the metabolic syndrome in children and adolescents. *Int J Pediatr Obes.* 2010;5(3):202-213.
37. Kelishadi R, Ardalan G, Gheiratmand R, Gouya MM, Razaghi EM, Delavari A. Association of physical activity and dietary behaviours in relation to the body mass index in a national sample of Iranian children and adolescents: CASPIAN Study. *Bulletin of the World Health Organization.* 2007;85(1):19-26.
38. Lawlor D, Riddoch C, Page A, Andersen LB, Wedderkopp N, Harro M. Infant feeding and components of the metabolic syndrome: findings from the European Youth Heart Study. *Arch Dis Childhood.* 2005;90(6):582-588.
39. Kries R, Koletzko B, Sauerwald T, Mutius E. Does Breast-Feeding Protect Against Childhood Obesity? Short & long term effects of breast feeding on child health. 2002:29-39.
40. Kelishadi R, Gouya MM, Adeli K, Ardalan G, Gheiratmand R, Majdzadeh R. Factors associated with the metabolic syndrome in a national sample of youths: CASPIAN Study. *Nutrition, Metabolism Cardiovasc Dis.* 2008;18(7):461-470.
41. Miles HL, Hofman PL, Cutfield WS. Fetal origins of adult disease: a paediatric perspective. *Revi Endocrine Metabolic Disorders.* 2005;6(4):261-268.
42. Phenekos C. Influence of fetal body weight on metabolic complications in adult life: review of the evidence. *J Pediatr Endocrinol Metabol.* 2001;14:1361.
43. Phillips D. Birth weight and the future development of diabetes. A review of the evidence. *Diab Care.* 1998;21:B150.
44. Savva S, Tornaritis M, Savva M, Kourides Y, Panagi A, Silikiotiou N. Waist circumference and waist-to-height ratio are better predictors of cardiovascular disease risk factors in children than body mass index. *Int J obesity and related metabolic disorders: J Int Assoc Study Obes.* 2000;24(11):1453.
45. Weiss R. Metabolic syndrome in childhood—causes and effects. *Endocr Dev.* 2010;19:62-72.
46. Goran MI, Gower BA. Relation between visceral fat and disease risk in children and adolescents. *Am J Clin Nutr.* 1999;70(1):149S-156S.
47. Ilyas M, Ellis E. Management of childhood hypertension: a guide for primary care physicians. *J Arkansas Med Soc.* 2006;103(6):137.
48. Salem Z. Prevalence Of Hypertension And Its Association With The Anthropometric Indices In Adolescent Girls In Rafsanjan, 2007. *J Rafsanjan Uni Med Sci Health Services.* 2010.
49. Kelishadi R, Heshmat R, Motlagh ME, Majdzadeh R, Keramatian K, Qorbani M. Methodology and early findings of the third survey of CASPIAN study: A national school-based surveillance of students' high risk behaviors. *Int J Prev Med* 2012;3:394-401.
50. Kelishadi R. Childhood overweight, obesity, and the metabolic syndrome in developing countries. *Epidemiol Rev.* 2007;29:62-76.